Title	Landscape Resilience Data			
Abstract	These data were collected as part of the <u>NSW Grazing Study</u> . This study investigates how past grazing pressure has influenced the resilience of all three vegetation communities. We assessed three key ecosystem processes or components that provide important information on the likely resilience of reserves following removal, or reduction in the densities of, domestic stock. These processes are 1) soil nutrient cycling, 2) soil hydrology and 3) seed bank dynamics.			
	For each community, we examined 18 individual sites scattered widely across the full range of the vegetation community, in three distinct groups ($n = 54$). In Cypress pine, the groups were selected along a gradient from north to south i.e. Cobar/Yathong (northern group), Griffith/Merriwagga (central group) and Narrandera/ Buckingbong (southern group). Similarly, Black box sites spanned a north-south gradient from Hillston (northern group), Griffith (central group) to Edwards River (southern group). River red gum sites were located along a gradient from east to west: Corowa (eastern group), Millewa (central group) and Deniliquin (western group). Each group contained three pairs of sites, with each pair corresponding to one of three different levels of grazing intensity; undisturbed (poor condition) by herbivore grazing. The assignment was based on attributes such as erosion extent, presence of exotic plants, groundstorey vegetation cover, the number of livestock and native herbivores (kangaroos, <i>Macropus spp</i>). For example, undisturbed (high condition) sites had an extensive cover of biocrusts, extensive litter cover, a plant community composition dominated by native species with a good mixture of annuals and perennials, abundant large grass butts, little evidence of erosion, high scores for the stability index and little evidence of grazing by livestock (based on dung counts and stock tracks). Highly disturbed sites intermediate between the two.			
	Floristic sampling - At each of the 54 sites we established a 100 m transect, perpendicular to the main watering point, along which we positioned three large quadrats (5 m x 5 m) at 0 m, 50 m and 100 m. Within these plots we centrally located a smaller (0.5 m x 0.5 m) quadrat. Within the large quadrats we conducted a full floristic survey (scoring the cover and abundance of all vascular plants).			
	Grazing Intensity - To assess recent grazing intensity, we identified and counted the dung or pellets of all herbivores (cattle, sheep/goat, kangaroo) within the large ($5m \times 5m$) and small ($0.5m \times 0.5m$) (kangaroo, rabbit, sheep/goat) quadrats. For cattle, the count of dung events from three $5m \times 5m$ quadrats. For sheep, goat, deer, kangaroo, rabbit the count of pellets from three $5m \times 5m$ quadrats and three $0.5m \times 0.5m$ quadrats.			
	Soil chemical and biological assessment - At each of the 0m, 50m and 100m positions along the site transect we collected two samples of the top 5 cm of soil with a soil corer. Samples were placed into one bag and a subsample of about 100 g taken for microbial and soil chemical analyses. Soil enzyme concentrations and soil nutrient pools were analysed at UNSW. Gene sequencing, using the Illumina MiSeq platform for bacteria and fungi, were carried out using the Next Generation Genome Sequencing Facility at the University of Western Sydney.			
	Soil infiltration measurements - We measured infiltration at the Black box and Cypress pine sites only, i.e. 36 of the 54 Stage III sites. At each microsite we used two disk permeameters, simultaneously, to measure sorptivity and steady-state infiltration under ponding (+ 10 mm) and under tension (- 40 mm). Disk permeameters were placed within 30 cm of each other, or as close as possible for measurements over grass butts. The tension permeameter was placed on a thin bed of sand to provide a uniform contact with the soil surface, and the ponded permeameter on a steel ring above a pond of water about 5 cm deep. For shrubs and trees, the permeameters were placed in the centre of the canopy. For grass microsites, the above-ground material was clipped and the permeameters placed directly over the grass butt. The permeameters were run for at least 12 minutes to ensure that they had achieved steady-state infiltration. This method allowed us to calculate values for both stages of infiltration: sorptivity, the early stage, and steady-state infiltration, the final stage of			

infiltration: sorptivity, the early stage, and steady-state infiltration, the final stage of infiltration. River red gum sites were excluded from soil infiltration measures because their heavy textured soils are prone to developing deep cracks, making the assessment of infiltration technically very difficult.

Seedbank dynamics - At each site (54) all three communities (Redgum, black box and cypress pine) two core samples were collected at 5 points to 5cm deep (0m, 50m, 100m, 150m 200m) at four patch types (tree, shrub, grass, open) along a 200m transect.

Seedling emergence experiments were conducted to determine the composition of the germinable soil seed bank (Thompson & Grime, 1979). The sieved soil was spread evenly (~5 mm deep) over sterilised sand in commercial germination trays (35 cm \times 14 cm) and placed in an unheated greenhouse. The trays were watered regularly to

keep the soil moist or at field capacity and the position of all trays was randomly allocated to account for a possible bias associated with tray position. Ten control trays, i.e. trays containing only sterilised sand, were evenly distributed in the greenhouse to control for glasshouse weeds and seeds within the sterilised sand. Emerging plants were counted and removed following identification,or representative samples repotted to grow on to confirm sample identification. The seedling emergent trial ran from spring 2016 to late autumn (242 days).

Resource locator

Data Quality	Name: Data Quality Statement			
<u>Statement</u>	Protocol: WWW:DOWNLOAD-1.0-httpdownload			
	Description:			
	Data quality statement for Plant Species Cover and Abundance			
	Function: download			
Landscape	Name: Landscape Resilience data			
<u>Resilience data</u>	Protocol: WWW:DOWNLOAD-1.0-httpdownload			
	Description:			
	Spreadsheet detailing location(s), plant cover abundance, soil surface condition, infiltration, dung, soil chemical assessment, microbes, enzymes and seedbank data.			
	Function: download			
Unique resource identifier				
Code	fe9648f9-3fbd-42d9-b88c-ed9e8caad655			
Presentation form	tableDigital			
Edition	Original (raw data)			
Dataset language	eng			
Metadata stan	dard			
Name	ANZLIC Metadata Profile: An Australian/New Zealand Profile of AS/NZS ISO 19115:2005, Geographic information - Metadata			
Version	1.1			
Dataset URI	https://datasets.seed.nsw.gov.au/dataset/fe9648f9-3fbd-42d9-b88c-ed9e8caad655			
Purpose	Decision support			
Status	completed			
Spatial reference system				
Authority code	GDA94 Geographic (Lat\Long)			
Code identifying the spatial reference system	4283			
Topic category Biota				

Keyword set	
keyword value	FLORA
	VEGETATION
	ECOLOGY
	SOIL
Originating controlled vocabulary	
Title	ANZLIC Search Words
Reference date	2008-05-16
Geographic location	
West bounding longitude	141.064453
East bounding longitude	148.139648
North bounding latitude	-35.995785
South bounding latitude	-31.802893
NSW Place Name	Central West, Riverina and Western NSW
Vertical extent information	
Minimum value	-100
Maximum value	2228
Coordinate reference system	
Authority code	urn:ogc:def:cs:EPSG::
Code identifying the coordinate reference system	5711
Temporal extent	
Begin position	2013-07-01
End position	N/A
Dataset reference date	
Date type	creation
Effective date	2013-07-01
Resource maintenance	
Maintenance and update frequency	notPlanned
Contact info	
Organisation name	Department of Planning, Industry and Environment
Full postal address	PO Box A290
	Sydney South
	NSW
	1232
	Australia
	data.broker@environment.nsw.gov.au

Telephone number			131555		
Facsimile nu	mber		02 9995 5999		
Email addres	SS		data.broker@environment.nsw.gov.au		
Responsible	party role		pointOfContact		
Constraint	set				
Use constraints	This data is provided under a Creative Commons Attribution 4.0 licence <u>http://creativecommons.org/licenses/by/4.0</u> Attribute 'Department of Planning, Industry and Environment ' in publications using this data.				
Limitations on public access					
Scope			dataset		
Responsible party					
Contact position		Data Broker			
Organisation name		Department of Planning, Industry and Environment			
Full postal address		PO Box A290			
		Sydney South			
		NSW			
		1232			
		Australia			
		data.broker@e	nvironment.nsw.gov.au		
Telephone number		131555			
Facsimile number		02 9995 5999			
Email address		data.broker@environment.nsw.gov.au			
Web address		http://www.planning.nsw.gov.au/			
Responsible party role		pointOfContact			

Metadata point of contact					
Contact position	Data Broker				
Organisation name	Department of Planning, Industry and Environment				
Full postal address	PO Box A290				
	Sydney South				
	NSW				
	1232				
	Australia				
	data.broker@environment.nsw.gov.au				
Telephone number	131555				
Facsimile number	02 9995 5999				
Email address	data.broker@environment.nsw.gov.au				
Responsible party role	distributor				
Metadata date	2013-07-01				
Metadata language	eng				